

N,O- vs N,C-Chelation in Half-Sandwich Iridium Complexes: A Dramatic Effect on Enantioselectivity in Asymmetric Transfer Hydrogenation of Ketones

Gang Zhou,^{†,‡} Ahmed H. Aboo,[‡] Craig M. Rebertson,[‡] Ruixia Liu,[†] Zhenhua Li,^{#,‡} Konstantin Luzyanin,[‡] Neil G. Berry,[‡] Weiping Chen,^{*,†} and Jianliang Xiao^{*,†,‡}

[†]School of Pharmacy, Fourth Military Medical University, Xi'an, 710032, China


[‡]Department of Chemistry, University of Liverpool, Liverpool, L69 7ZD, United Kingdom

[#]Department of Chemistry, Fudan University, Shanghai, 200438, China

S Supporting Information

ABSTRACT: Cyclometalation of $[\text{Cp}^*\text{IrCl}_2]_2$ with methyl (S)-2-phenyl-4,5-dihydrooxazole-4-carboxylate in the presence of NaOAc selectively led to a N,C- or N,O-chelated $\text{Cp}^*\text{Ir(III)}$ complex, depending on whether or not water was present in the reaction. While derived from the same precursor, these two complexes behaved in a dramatically different manner in asymmetric transfer hydrogenation (ATH) of ketones by formic acid, with the N,O-chelated complex being much more selective and active. The sense of asymmetric induction is also different, with the N,O-complex affording S while the N,C-analogue R alcohols. Further study revealed that the nature of the base additive considerably impacts the enantioselectivity and the effective $\text{HCOOH}/\text{amine}$ ratios. These observations show the importance of ligand coordination mode and using the right base for ATH reactions.

KEYWORDS: N,O-chelation, N,C-chelation, half-sandwich iridium complexes, cyclometalation, asymmetric transfer hydrogenation



N,C-Chelated half-sandwich iridium complexes of type 1 have received a great deal of attention in the past decade, finding numerous applications in catalysis among others (Figure 1).¹ In 2008, Ikariya and co-workers reported that




Figure 1. Selected examples of N,C- and N,O-chelated half-sandwich iridium complexes.

complex 2 catalyzes the aerobic oxidation of alcohols.² When the metalacycle was made chiral with a simple chiral amine,

oxidative kinetic resolution of racemic alcohols was shown to be feasible. In the same year, Pfeffer, Janssen, Feringa, de Vries et al. found that complex 3 with a simple amine ligand is a good catalyst for racemization of alcohols.³ In 2009, Crabtree and co-workers disclosed complex 4 with 2-phenylpyridine as a ligand for water oxidation.⁴ In 2010, one of our groups demonstrated that the ketimine-ligated complexes 5 are powerful catalysts for the reductive amination of a wide variety of carbonyl compounds.⁵ The following years have witnessed flourishing applications of half-sandwich cyclometalated iridium complexes in catalysis, including hydrogenation, reductive amination, dehydrogenation, oxidation, alkylation, racemisation, hydrosilylation, hydroamination, polymerization, and related reactions.^{1,6}

The somewhat related N,O-chelated half-sandwich complexes of iridium derived from α - and β -amino acids, 2-pyridylacetic acid, picolinic acid, or even peptide ligands have been known for decades.⁷ However, they have only scarcely been used in catalysis. Examples are found in the α -amino-acid-derived N,O-chelated complex 6, which catalyzes the asymmetric transfer hydrogenation (ATH) of ketones⁷ⁿ

Received: May 28, 2018

Revised: July 26, 2018

Published: July 30, 2018

51 and complex **7** as a highly reactive and selective catalyst for the
52 alkylation of amines with alcohols.^{7p} The iridium complex **8**
53 bearing a 2-pyridylacetic-acid-derived ligand is an efficient cata-
54 lyst for the dehydrogenation of alcohols.^{7o}

55 In continuing our exploration of N,C-chelated iridium com-
56 plexes in catalysis,^{1d} we targeted a simple chiral complex **9**,
57 anticipating that it might enable asymmetric reduction of
58 imines. The imino substrate could be activated by the carbox-
59 ylic acid (R = H) or the ester (R = alkyl) via hydrogen bonding
60 and thereby positioned, facilitating enantioselective hydride
61 transfer as illustrated in Figure 2.⁸ However, the outcome of




Figure 2. Target catalyst and proposed mode of asymmetric reduction of imines involving secondary interactions.

62 our endeavor is totally unexpected. The oxazoline ligand was
63 found to form, surprisingly, either a N,C- or a N,O-chelated half-
64 sandwich Ir(III)-complex and remarkably, this mode of chelation
65 has a dramatic effect on the enantioselectivity of the Cp*Ir(III)
66 complex-catalyzed ATH of ketones. While both N,C- and N,O-
67 chelated half-sandwich complexes have been well-documented
68 in the literature, little is known of how the difference in the
69 coordination mode of the ligand may affect their catalytic activ-
70 ity and selectivity.

71 Cyclometalation through C–H activation is a well-established
72 method for the synthesis of transition-metal complexes bearing
73 η^2 -C₂X (X = C, N, O) ligands.¹ According to a general procedure
74 for the preparation of cyclometalated complexes,^{5,9} methyl
75 (S)-2-phenyl-4,5-dihydrooxazole-4-carboxylate **10** was reacted
76 with [Cp*IrCl₂]₂ at room temperature in the presence of anhy-
77 drous NaOAc. The reaction afforded a mixture of two half-
78 sandwich Cp*Ir(III) complexes: the expected N,C-chelated
79 complex **11a** and an “abnormal” N,O-chelated complex **11b**, in
80 a ratio of **11a**:**11b** = 1:3.5 (Table 1, entry 1). Delightfully, the
81 ratio of **11a** to **11b** was found to be variable with the amount
82 of water in the solvent. Thus, when CH₂Cl₂ dried over CaH₂
83 was used, the ratio of **11a** increased with **11a**:**11b** = 1:1 (entry 2,
84 Table 1), and introducing 4 Å molecular sieves to this reaction
85 afforded the N,C-chelated complex **11a** as the sole product
86 (entry 3, Table 1). In sharp contrast, using wet CH₂Cl₂ led to
87 the exclusive formation of the N,O-chelated complex **11b**
88 (entry 4, Table 1). Most likely, **11b** is formed via initial coordi-
89 nation of the ester moiety to the Lewis acidic Ir(III) center,
90 followed by hydrolysis with water, as illustrated in Table 1.
91 In the absence of an ester group, cyclometalation occurs with
92 or without water (See Section 8 in the Supporting Information
93 (SI).) Both **11a** and **11b** are air-stable complexes. Attempts to
94 convert one to the other under various conditions, e.g., by
95 adding an acid or a base or raising the temperature, have not
96 been successful. The structures of **11a** and **11b** were
97 determined by single-crystal X-ray diffraction and are shown
98 in Figure 3.

99 Pure **11b** exists in solution as a mixture of two diastereomers
100 (ratio of 9.8:1), because of the presence of chiral centers at
101 iridium and the ligand. ¹H NMR monitoring of the freshly pre-
102 pared solution of **11b** in dry CDCl₃ or CD₃OD in the –50 °C
103 to +40 °C range indicated that the diastereomeric ratio does not

Table 1. Synthesis of Cyclometalated Cp*Ir(III) Complexes **11a** and **11b**^a

entry	solvent	additive	yield ^b (%)	11a : 11b ^c
1	CH ₂ Cl ₂ ^d	no	94	1:3.5
2	dried CH ₂ Cl ₂ ^e	no	92	1:1
3	dried CH ₂ Cl ₂ ^e	4 Å MS (50 mg/mL)	89	>99:1
4	CH ₂ Cl ₂ ^d	H ₂ O (2%, v/v)	97	<1:99

^aConditions: ligand (0.49 mmol), [Cp*IrCl₂]₂ (0.22 mmol), NaOAc (4.9 mmol), DCM (10 mL), rt, 24 h. ^bIsolated yield. ^cProduct ratio determined by ¹H NMR of the crude reaction mixture. ^dUsed as received. ^eDried over CaH₂.




Figure 3. Molecular structures of **11a** and **11b** determined by single-crystal X-ray diffraction (XRD). For **11a**, the selected bond distances are as follows: Ir1–Cl1, 2.4138(10) Å; Ir1–N1, 2.078(5) Å; Ir1–C1, 2.056(6) Å; Ir1–avgC(Cp*), 2.189(15) Å. For **11a**, the selected bond angles are as follows: N1–Ir1–Cl1, 87.47(15)°; C1–Ir1–Cl1, 86.84(18)°; C1–Ir1–N1, 77.7(3)°. For **11b** (solvent omitted for clarity), the selected bond distances (Å) are as follows: Ir1–Cl1, 2.404(2) Å; Ir1–O3, 2.152(7) Å; Ir1–N1, 2.092(8) Å; Ir1–avgC(Cp*), 2.142(23) Å. For **11b** (solvent omitted for clarity), the selected bond angles (deg) are as follows: O3–Ir1–Cl1, 83.6(2)°; N1–Ir1–Cl1, 88.3(3)°; N1–Ir1–O3, 77.0(2)°.

change noticeably by varying the temperature or solvent, even
after 24 h. No changes in the diastereomeric ratio were also
observed upon the addition of [Bu₄N]Br or [Bu₄N]I (5 equivs).
Prolonged heating of the mixture with [Bu₄N]Br or [Bu₄N]I
(40 °C, longer than 1 h) resulted in the gradual change of the
solution color from orange to red, presumably indicating the
replacement of the chloride with Br or I. The addition of an
excess amount of acetic acid (5 equivs) or a mixture of acetic
acid and isopropylamine did not alter the structure of **11b** or
its diastereomeric ratio either. Similarly, **11a** appears as a mix-
ture of two diastereomers, the ratio of which, however, is consid-
erably higher (>20:1), and addition of acetic acid and isopro-
pylamine to a solution of **11a** in CDCl₃ brought about no
notable effect, as shown by ¹H NMR (see Section 7 in the SI).

The fact that **11a** and **11b** differs mainly in the coordination
mode of the chiral ligand prompted us to compare their ability

Table 2. Comparison of ATH of *p*-Nitroacetophenone under Various Conditions^a

entry	catalyst	solvent	time (h)	conversion ^b (%)	enantiomeric excess, ee ^c (%)
1	11a	CH ₂ Cl ₂	15	75	4 (R)
2	11b	CH ₂ Cl ₂	2	100	73 (S)
3 ^d	11a	F/T	15	61	2 (R)
4 ^d	11b	F/T	15	96	38 (S)
5	11a	MeOH	15	80	4 (R)
6	11b	MeOH	15	97	53 (S)
7	11a	<i>i</i> PrOH	15	71	2 (R)
8	11b	<i>i</i> PrOH	15	99	40 (S)
9	11a	toluene	15	42	2 (R)
10	11b	toluene	15	100	42 (S)
11	11a	H ₂ O	15	54	3 (R)
12	11b	H ₂ O	15	85	27 (S)
13 ^e	11a	aq. solution of HCO ₂ H/HCO ₂ Na	15	58	0
14 ^e	11b	aq. solution of HCO ₂ H/HCO ₂ Na	15	100	37 (S)

^aConditions: substrate (0.2 mmol), catalyst (0.002 mmol), azeotropic F/T solution (0.5 mL), solvent (2 mL), room temperature. ^bDetermined by ¹H NMR of the crude reaction mixture. ^cDetermined by HPLC. ^dAzeotropic F/T solution (2.5 mL) was used with no additional solvent. ^eAqueous formate solution used (pH 4.5).

of catalyzing ATH reactions.¹⁰ First, we tested the catalytic performance of **11a** and **11b** in the ATH of ketones, choosing the reduction of *p*-nitroacetophenone as a model reaction. As can be seen from Table 2, in the presence of 1% of **11a** or **11b**, *p*-nitroacetophenone could be reduced by using an azeotropic mixture of formic acid/triethylamine (F/T) in CH₂Cl₂ at room temperature. However, the outcome is remarkably different. Thus, while the N,C-chelated **11a** showed a very low catalytic activity (75% conversion in 15 h) and extremely low enantioselectivity (4% ee), the N,O-analogue **11b** was much more active and enantioselective (100% conversion in 2 h, 73% ee). Of further notice is that the configuration of the products obtained with **11a** and **11b** is opposite. This sharp difference was repeated in other solvents as well, reinforcing the contrast brought about by a simple change in ligand coordination mode and the superiority of the N,O-chelated **11b** (Table 2, entries 5–14). The best enantioselectivity was observed in CH₂Cl₂ with **11b**. These observations suggest that, although **11a** and **11b** bear chiral ligands of similar original structure, the differing coordination mode of the ligands impacts the mechanism of how they affect the ATH and, particularly, the step of hydride transfer, where the enantioselectivity is likely to be determined. Bearing in mind that the ratio of F/T may affect the enantioselectivity of ATH of ketones,¹¹ we also examined the effect of this parameter on the ATH with the more effective catalyst **11b**. As shown in Figure 4, the F/T ratio indeed impacts on the enantiomeric excess (ee) of the ATH in question, with the highest ee observed in a narrow window of ca. 2.5–3. More interestingly, variation of the nature of the amine used brought about a hitherto little-noticed finding, i.e., both the nature of the amine and its ratio with HCOOH considerably affect the enantioselectivity of the ATH. Among the tested amines, the HCO₂H-*i*-PrNH₂ (2:1) mixture gave the highest enantioselectivity, with a significantly widened window of effective HCOOH/amine ratios. While the reason for the varying effect of amines on the ee is not entirely clear at the moment, the observation calls for attention when examining other catalysts for ATH reactions with formic acid, where NEt₃ has been used as a base almost exclusively in the past decades.^{10a–d,g–n}




Figure 4. Effect of amines and the molar ratio of HCOOH/amine on the enantioselectivity of the ATH with catalyst **11b**. Conditions: *p*-nitroacetophenone (0.2 mmol), catalyst (0.002 mmol), HCOOH/amine solution (0.5 mL), DCM (2 mL), room temperature. The ee value was determined by high-performance liquid chromatography (HPLC).

Under the optimized conditions, we made further comparison of **11a** with **11b** in the ATH of acetophenones bearing either electron-donating or electron-withdrawing substituents on the aromatic (Table 3). As with the reduction using an azeotropic mixture of F/T as hydrogen source, the **11b**-catalyzed ATH of all four tested acetophenones with the HCO₂H-*i*-PrNH₂ (2:1) mixture gave excellent enantioselectivity in each case (see Table 3, entries 2, 4, 6, 8, and 10), while the performance of **11a** was much poorer (Table 3, entries 1, 3, 5, 7, and 9). These observations substantiate further the assertion that the coordination mode of ligands can exert a significant effect on the activity and enantioselectivity of ATH reactions.

The scope of substrates was subsequently examined with complex **11b** using the HCO₂H-*i*-PrNH₂ (2:1) mixture

Table 3. Comparison of ATH of Aromatic Ketones Catalyzed by 11a and 11b^a

entry	R	catalyst	conversion ^b (%)	enantiomeric excess, ee ^c (%)
1	H	11a	6	4 (R)
2	H	11b	23	98 (S)
3	<i>o</i> -OMe	11a	8	6 (R)
4	<i>o</i> -OMe	11b	20	93 (S)
5	<i>p</i> -OMe	11a	10	7 (R)
6	<i>p</i> -OMe	11b	23	92 (S)
7	<i>p</i> -Br	11a	20	3(R)
8	<i>p</i> -Br	11b	65	99 (S)
9	<i>p</i> -NO ₂	11a	30	5 (R)
10	<i>p</i> -NO ₂	11b	100	93 (S)

^aConditions: substrate (0.2 mmol), catalyst (0.002 mmol), HCOOH/amine (2:1) solution (0.5 mL), DCM (2 mL), room temperature, 3 h. ^bDetermined by ¹H NMR of the crude reaction mixture. ^cDetermined by HPLC.

173 as a hydrogen source (see [Figure 5](#)). All aromatic ketones
174 could be reduced with excellent enantioselectivities (90%–




Figure 5. ATH of various aromatic ketones with complex **11b**. Isolated yields are given. For more details, see the [SI](#).

175 99% ee). However, the catalyst shows a low activity toward
176 acetophenones that bear highly electron-donating substituents
177 or sterically more demanding ones, e.g., 4-hydroxyacetophenone
178 and α -substituted acetophenones. We note that
179 electron-rich ketones have been challenging for ATH catalysts
180 in general, and only a few examples of ATH of hydroxyacetophenones
181 are known.¹² Still disappointingly, neither **11b** nor **11a** was found to
182 be enantioselective in the ATH of imines.
183

A plausible mechanism for the **11b**-catalyzed ATH is shown in **Figure 6**. The steps leading to the iridium-hydride from **11b**




Figure 6. Suggested mechanism for the ATH of ketones with the N,O-chelated iridium complex. The ammonium cation may hydrogen bond with the N,O-ligand throughout the catalytic cycle. The suggested transition state of hydride transfer is supported by a DFT calculation. (Ar = Ph. For details, see Section 13 of the SI.)

would be expected to be similar to those proposed for the N,C-chelated iridacycles.¹³ It is the hydride transfer step that sets this catalyst apart from other N,O- or N,C-chelated iridium catalysts. We hypothesize that the ammonium cation participates in the transition state of this enantioselectivity-determining step, hydrogen-bonding both the N,O-ligand via its carboxylate oxygen and the ketone substrate through its carbonyl oxygen. Such a hydrogen bonding network would be expected to lower the barrier of the transition state and enhance the enantioselectivity of the hydride transfer. Density functional theory (DFT) modeling of the hydride-transfer step revealed that the isopropylammonium cation can indeed participate in the transition state and further showed, consistent with the experiment, in that it is the S alcohol that is to be favored ($\Delta\Delta G^\ddagger = 1.8$ kcal/mol). As shown in Figure 6, the transition state of the hydride transfer involves two protons of the ammonium cation strongly hydrogen-bonding with the oxygen atom of the carboxylate ligand (a; O...H distance = 1.92 Å)

and the acetophenone oxygen (c; O...H distance = 1.25 Å, indicating significant O–H bond formation) simultaneously. There also appear to be weaker interactions between these two protons and the ligand oxygen (b; 2.91 and 2.90 Å, respectively) (see Section 13 of the SI for more details). The existence of the hydrogen bonding in question may not be unexpected, as ammonium cations are widely known to form moderately strong hydrogen bonds with various carbonyl compounds.¹⁴ In ATH reactions, ligand-induced hydrogen bonding has been well-established since the pioneering work of Noyori and Hashiguchi;¹⁵ however, examples of hydrogen bonding enabled by carboxylate ligands are relatively rare.¹⁶ The calculated transition state in Figure 6 also indicates why the nature of the ammonium cation affects significantly the enantioselectivity, with the cation directly involved in the enantioselectivity-determining step. What remains to be delineated is how the other cations, e.g., Et₃NH⁺, participate in the transition state and thereby affect the ee, although primary ammonium cations appear to form stronger hydrogen bonds with ketones than tertiary ones.¹⁴

In summary, we have demonstrated that (1) a N,C- or a N,O-chelated half-sandwich Cp*Ir(III)-complex can be selectively prepared from the reaction of methyl (S)-2-phenyl-4,5-dihydrooxazole-4-carboxylate with [Cp*IrCl₂]₂ by simply changing the reaction conditions; (2) the mode of chelation has a dramatic effect on the enantioselectivity of the Cp*Ir(III) complex-catalyzed ATH of ketones; and (3) the nature of the amine and its ratio with HCOOH significantly affect the enantioselectivity of the N,O-complex-catalyzed ATH reaction.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscatal.8b02068.

- Experimental procedures and characterization data, ¹H and ¹³C NMR spectra (PDF)
- Crystallographic information for C₂₁H₂₅ClIrNO₃ (CIF)
- Crystallographic information for C₂₁H₂₅Cl₃IrNO₃ (CIF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: wpchen@fmmu.edu.cn.

*E-mail: jxiao@liv.ac.uk.

ORCID

Zhenhua Li: 0000-0002-5636-9865

Jianliang Xiao: 0000-0003-2010-247X

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China (21272271). We are grateful to China Scholarship Council (File No.201503170380) for funding and the University of Liverpool for support.

■ REFERENCES

- (1) Recent reviews, see: (a) Liu, J.; Wu, X.; Iggo, J. A.; Xiao, J. Half-sandwich Iridium Complexes—Synthesis and Applications in Catalysis. *Coord. Chem. Rev.* **2008**, *252*, 782–809. (b) Han, Y.-F.; Jin, G.-X. Cyclometalated [Cp* M (C[^]X)] (M = Ir, Rh; X = N, C, O, P) Complexes. *Chem. Soc. Rev.* **2014**, *43*, 2799–2823. (c) Michon, C.;

- MacIntyre, K.; Corre, Y.; Agbossou-Niedercorn, F. Pentamethylcyclopentadienyl Iridium(III) Metallocycles Applied to Homogeneous Catalysis for Fine Chemical Synthesis. *ChemCatChem* **2016**, *8*, 1755–1762. (d) Wang, C.; Xiao, J. Iridacycles for Hydrogenation and Dehydrogenation reactions. *Chem. Commun.* **2017**, *53*, 3399–3411.
- (2) (a) Arita, S.; Koike, T.; Kayaki, Y.; Ikariya, T. Aerobic Oxidative Kinetic Resolution of Racemic Secondary Alcohols with Chiral Bifunctional Amido Complexes. *Angew. Chem., Int. Ed.* **2008**, *47*, 2447–2449. (b) Arita, S.; Koike, T.; Kayaki, Y.; Ikariya, T. Aerobic Oxidation of Alcohols with Bifunctional Transition-Metal Catalysts Bearing C–N Chelate Ligands. *Chem.—Asian J.* **2008**, *3*, 1479. (c) Arita, S.; Koike, T.; Kayaki, Y.; Ikariya, T. Synthesis and Reactivities of Cp* Ir Amide and Hydride Complexes Bearing C–N Chelate Ligands. *Organometallics* **2008**, *27*, 2795–2802.
- (3) (a) Haak, R. M.; Berthiol, F.; Jerphagnon, T.; Gayet, A. J. A.; Tarabiono, C.; Postema, C. P.; Ritleng, V.; Pfeffer, M.; Janssen, D. B.; Minnaard, A. J.; Feringa, B. L.; de Vries, J. G. Dynamic Kinetic Resolution of Racemic Beta-haloalcohols: Direct Access to Enantioenriched Epoxides. *J. Am. Chem. Soc.* **2008**, *130*, 13508–13509. (b) Jerphagnon, T.; Gayet, A. J. A.; Berthiol, F.; Ritleng, V.; Mršić, N.; Meetsma, A.; Pfeffer, M.; Minnaard, A. J.; Feringa, B. L.; de Vries, J. G. Fast Racemisation of Chiral Amines and Alcohols by Using Cationic Half-Sandwich Ruthenium- and Iridacycle Catalysts. *Chem.—Eur. J.* **2009**, *15*, 12780–12790.
- (4) Hull, J. F.; Balcells, D.; Blakemore, J. D.; Incarvito, C. D.; Eisenstein, O.; Brudvig, G. W.; Crabtree, R. H. Highly Active and Robust Cp* Iridium Complexes for Catalytic Water Oxidation. *J. Am. Chem. Soc.* **2009**, *131*, 8730–8731.
- (5) Wang, C.; Pettman, A.; Basca, J.; Xiao, J. A Versatile Catalyst for Reductive Amination by Transfer Hydrogenation. *Angew. Chem., Int. Ed.* **2010**, *49*, 7548–7552.
- (6) For recent examples of the N,C-chelated half-sandwich Ir-complexes catalyzed reactions, see: (a) Kim, H.; Chang, S. Selective Heterogeneous CO₂ Electroreduction to Methanol. *ACS Catal.* **2015**, *5*, 6665–6669. (b) Arthurs, R. A.; Ismail, M.; Prior, C. C.; Oganessian, V. S.; Horton, P. N.; Coles, S. J.; Richards, C. J. Enantiopure Ferrocene-Based Planar-Chiral Iridacycles: Stereospecific Control of Iridium-Centred Chirality. *Chem.—Eur. J.* **2016**, *22*, 3065–3072. (c) Arthurs, R. A.; Horton, P. N.; Coles, S. J.; Richards, C. J. Phenyl vs. Ferrocenyl Cyclometallation Selectivity: Diastereoselective Synthesis of an Enantiopure Iridacycle. *Eur. J. Inorg. Chem.* **2017**, *2017*, 229–232. (d) Sato, Y.; Kayaki, Y.; Ikariya, T. Comparative Study of Bifunctional Mononuclear and Dinuclear Amidoiridium Complexes with Chiral C–N Chelating Ligands for the Asymmetric Transfer Hydrogenation of Ketones. *Chem.—Asian J.* **2016**, *11*, 2924–2931. (e) Semwal, S.; Mukkatt, I.; Thenarukandiyil, R.; Choudhury, J. Small “Yaw” Angles, Large “Bite” Angles and an Electron-Rich Metal: Revealing a Stereoelectronic Synergy to Enhance Hydride-Transfer Activity. *Chem.—Eur. J.* **2017**, *23*, 13051–13057. (f) Fujita, K.; Tamura, R.; Tanaka, Y.; Yoshida, M.; Onoda, M.; Yamaguchi, R. Dehydrogenative Oxidation of Alcohols in Aqueous Media Catalyzed by a Water-Soluble Dicationic Iridium Complex Bearing a Functional N-Heterocyclic Carbene Ligand without Using Base. *ACS Catal.* **2017**, *7*, 7226–7230. (g) Matsunami, A.; Kuwata, S.; Kayaki, Y. Hydrodefluorination of Fluoroarenes Using Hydrogen Transfer Catalysts with a Bifunctional Iridium/NH Moiety. *ACS Catal.* **2016**, *6*, 5181–5185. (h) Corre, Y.; Werlé, C.; Brelot-Karmazin, L.; Djukic, J.-P.; Agbossou-Niedercorn, F.; Michon, C. Regioselective Hydro-silylation of Terminal Alkynes Using Pentamethylcyclopentadienyl Iridium(III) Metallocycle Catalysts. *J. Mol. Catal. A: Chem.* **2016**, *423*, 256–263. (i) Yao, Z.-J.; Li, K.; Li, P.; Deng, W. Mononuclear Half-sandwich Iridium and Rhodium Complexes through C–H Activation: Synthesis, Characterization and Catalytic Activity. *J. Organomet. Chem.* **2017**, *846*, 208–216. (j) Sato, Y.; Kayaki, Y.; Ikariya, T. Cationic Iridium and Rhodium Complexes with C–N Chelating Primary Benzylic Amine Ligands as Potent Catalysts for Hydrogenation of Unsaturated Carbon–Nitrogen Bonds. *Organometallics* **2016**, *35*, 1257–1264. (k) Petronilho, A.; Vivancos, A.; Albrecht, M. Ether Formation through Reductive Coupling of Ketones or

- Aldehydes Catalyzed by a Mesoionic Carbene Iridium Complex. *Catal. Sci. Technol.* **2017**, *7*, 5766–5774.
- (7) (a) Koch, D.; Sünkel, K.; Beck, W. Z. Metallkomplexe mit biologisch wichtigen Liganden. CLIV [1] Halbsandwich-Komplexe von Rhodium, Iridium, Ruthenium und phosphanhaltige Palladium- und Platin-Komplexe mit Sarkosinat, N-Methylalaninat und Ethylendiamin-*N,N'*-Diacetat. *Z. Anorg. Allg. Chem.* **2003**, *629*, 1322–1328. (b) Carmona, D.; Pilar Lamata, M.; Viguri, F.; San José, E.; Mendoza, A.; Lahoz, F. J.; García-Orduna, P.; Atencio, R.; Oro, L. A. N-Benzyl and N-Aryl Bis(Phospha-Mannich adducts): Synthesis and Catalytic Activity of the Related Bidentate Chelate Platinum Complexes in Hydroformylation. *J. Organomet. Chem.* **2012**, *717*, 152–163. (c) Jimineo, M. L.; Elguero, J.; Carmona, D.; Pilar Lamata, M.; San Jose, E. ¹H NMR Study of the Conformation of Metallapentacycles N-C-C-O-M [M = Rh(III) and Ir(III)] Resulting in a Karplus-Type Relationship for Vicinal H–C(sp³)–N(sp¹³)–H Coupling Constants. *Magn. Reson. Chem.* **1996**, *34*, 42–46. (d) Poth, T.; Paulus, H.; Elias, H.; Dücker-Benfer, C.; van Eldik, R. Kinetics and Mechanism of Water Substitution at Half-Sandwich Iridium(III) Aqua Cations Cp*Ir(A–B)(H₂O)^{2+/+} in Aqueous Solution (Cp* = η⁵-Pentamethylcyclopentadienyl Anion; A–B = Bidentate N,N or N,O Ligand). *Eur. J. Inorg. Chem.* **2001**, *2001*, 1361–1369. (e) Carmona, D.; Vega, C.; Lahoz, F. J.; Atencio, R.; Oro, L. A.; Pilar Lamata, M.; Viguri, F.; San Jose, E. Synthesis and Stereochemistry of Half-Sandwich Alkynyl Amino Acidate Complexes of Rhodium(III), Iridium(III), and Ruthenium(II). *Organometallics* **2000**, *19*, 2273–2280. (f) Carmona, D.; Lahoz, F. J.; Atencio, R.; Oro, L. A.; Lamata, M. P.; San José, E. Chiral Iridium(III) α-Amino Acidato Complexes (R₁Sn₂SC[–]) and (S₁Sn₂SC[–])-[(η⁵-C₅Me₅)Ir(L–proline)(C C CMe₃)]. *Tetrahedron: Asymmetry* **1993**, *4*, 1425–1428. (g) Grotjahn, D. B.; Groy, T. L. Formation and Structure of Coordinatively Unsaturated CpIr-Amino Acid Complexes. Kinetic and Thermodynamic Control in Highly Diastereoselective Complexation Reactions. *Organometallics* **1995**, *14*, 3669–3682. (h) Grotjahn, D. B.; Groy, T. L. Formation of Coordinatively Unsaturated Cp*Ir-Amino Acid Complexes and Their Highly Diastereoselective Complexation Reactions. *J. Am. Chem. Soc.* **1994**, *116*, 6969–6970. (i) Carmona, D.; Mendoza, A.; Lahoz, F. J.; Oro, L. A.; Lamata, M. P.; San José, E. Optically Active Pseudooctahedral Rhodium(III), Iridium(III), and Ruthenium(II) Complexes with α-Amino Acidato Ligands. Crystal structures of R₁SC₂N[–] and S₁SC₂N[–][(C₅Me₅)Ir(pro)-Cl]·12H₂O (Hpro = l-proline). *J. Organomet. Chem.* **1990**, *396*, C17–C21. (j) Grotjahn, D. B.; Joubert, C.; Hubbard, J. L. Highly Stereoselective Formation of Cp*IrCl Complexes of *N,N*-Dimethylamino Acids. *Organometallics* **1996**, *15*, 1230–1235. (k) Kramer, R.; Polborn, K.; Wanjek, H.; Zahn, I.; Beck, W. Chirale Halbsandwich-Komplexe von Rhodium(III), Iridium(III), Iridium(I) und Ruthenium(II) mit α-Aminosäure-Anionen. *Chem. Ber.* **1990**, *123*, 767–778. (l) Koch, D.; Hoffmüller, W.; Polborn, K.; Beck, W. Z. Metal Complexes of Biologically Important Ligands, CXXXVII [1]. Halbsandwich Complexes with N,O-Chelates and Schiff Bases of β-Amino Acids. *Z. Naturforsch., B: J. Chem. Sci.* **2001**, *56b*, 403–410. (m) Hoffmüller, W.; Dialer, H.; Beck, W. Z. Metal Complexes with Biologically Important Ligands, CLXI. Halbsandwich Complexes with tert-Leucine, Dipeptides, Pentaglycine and Glutathione. *Z. Naturforsch., B: J. Chem. Sci.* **2005**, *60*, 1278–1286. (n) Carmona, D.; Lahoz, F. J.; Atencio, R.; Oro, L. A.; Lamata, M. P.; Viguri, F.; San José, E.; Vega, C.; Reyes, J.; Joó, F.; Kathó, A. Trimerisation of the Cationic Fragments [(η-ring)M(Aa)]⁺[(η-ring) M = (η⁵-C₅Me₅)Rh, (η⁵-C₅Me₅)Ir, (η⁶-p-MeC₆H₄Pr)Ru; Aa = α-amino acidate) with Chiral Self-Recognition: Synthesis, Characterisation, Solution Studies and Catalytic Reactions of the Trimers [{(η-ring)M(Aa)}₃](BF₄)₃. *Chem.—Eur. J.* **1999**, *5*, 1544–1564. (o) Royer, A. M.; Rauchfuss, T. B.; Wilson, S. R. Coordination Chemistry of a Model for the GP Cofactor in the Hmd Hydrogenase: Hydrogen-bonding and Hydrogen-transfer Catalysis. *Inorg. Chem.* **2008**, *47*, 395–397. (p) Wetzlar, A.; Wöckel, S.; Schelwies, M.; Brinks, M. K.; Rominger, F.; Hofmann, P.; Limbach, M. Selective Alkylation of Amines with Alcohols by Cp* - Iridium(III) Half-Sandwich Complexes. *Org. Lett.* **2013**, *15*, 266–269.
- (8) For recent examples of secondary interactions between the amino groups and carboxylic acids in asymmetric hydrogenation, see: (a) Chen, W.; McCormack, P. J.; Mohammed, K.; Mbafor, W.; Roberts, S. M.; Whittall, J. Stereoselective Synthesis of Ferrocene-Based C₂-Symmetric Diphosphine Ligands: Application to the Highly Enantioselective Hydrogenation of α-Substituted Cinnamic Acids. *Angew. Chem., Int. Ed.* **2007**, *46*, 4141–4144. (b) Chen, W.; Spindler, F.; Pugin, B.; Nettekoven, U. ChenPhos: Highly Modular P-Stereogenic C₁-Symmetric Diphosphine Ligands for the Efficient Asymmetric Hydrogenation of α-Substituted Cinnamic Acids. *Angew. Chem., Int. Ed.* **2013**, *52*, 8652–8656. (c) Yao, L.; Wen, J.; Liu, S.; Tan, R.; Wood, N. M.; Chen, W.; Zhang, S.; Zhang, X. Highly Enantioselective Hydrogenation of α-Oxy Functionalized α,β-Unsaturated Acids Catalyzed by a ChenPhos-Rh complex in CF₃CH₂OH. *Chem. Commun.* **2016**, *52*, 2273–2276. (d) Chen, C.; Wang, H.; Zhang, Z.; Jin, S.; Wen, S.; Ji, J.; Chung, L. W.; Dong, X.-Q.; Zhang, X. Ferrocenyl Chiral Bisphosphorus Ligands for Highly Enantioselective Asymmetric Hydrogenation via Noncovalent Ion Pair Interaction. *Chem. Sci.* **2016**, *7*, 6669–6673. (e) Yin, X.; Chen, C.; Dong, X.-Q.; Zhang, X. Rh/Wudaphos-Catalyzed Asymmetric Hydrogenation of Sodium α-Arylethylsulfonates: A Method to Access Chiral α-Arylethylsulfonic Acids. *Org. Lett.* **2017**, *19*, 2678–2681. (f) Chen, C.; Zhang, Z.; Jin, S.; Fan, X.; Geng, M.; Zhou, Y.; Wen, S.; Wang, X.; Chung, L. W.; Dong, X.-Q.; Zhang, X. Enzyme-Inspired Chiral Secondary-Phosphine-Oxide Ligand with Dual Noncovalent Interactions for Asymmetric Hydrogenation. *Angew. Chem., Int. Ed.* **2017**, *56*, 6808–6812. (g) Yin, X.; Chen, C.; Li, X.; Dong, X.-Q.; Zhang, X. Rh/SPO-WudaPhos-Catalyzed Asymmetric Hydrogenation of α-Substituted Ethenylphosphonic Acids via Noncovalent Ion-Pair Interaction. *Org. Lett.* **2017**, *19*, 4375–4378. (h) Chen, C.; Wen, S.; Dong, X.-Q.; Zhang, X. Highly Stereoselective Synthesis and Application of P-chiral Ferrocenyl Bisphosphorus Ligands for Asymmetric Hydrogenation. *Org. Chem. Front.* **2017**, *4*, 2034–2038. (i) Chen, C.; Wen, S.; Geng, M.; Jin, S.; Zhang, Z.; Dong, X.-Q.; Zhang, X. A New Ferrocenyl Bisphosphorus Ligand for the Asymmetric Hydrogenation of α-methylene-γ-keto-carboxylic Acids. *Chem. Commun.* **2017**, *53*, 9785–9788. (j) Wen, S.; Chen, C.; Du, S.; Zhang, Z.; Huang, Y.; Han, Z.; Dong, X.-Q.; Zhang, X. Highly Enantioselective Asymmetric Hydrogenation of Carboxy-Directed α,α-Disubstituted Terminal Olefins via the Ion Pair Noncovalent Interaction. *Org. Lett.* **2017**, *19*, 6474–6477.
- (9) (a) Davies, D. L.; Al-Duaj, O.; Fawcett, J.; Giardiello, M.; Hilton, S. T.; Russell, D. R. Room-temperature Cyclometallation of Amines, Imines and Oxazolines with MCl Cp M Rh, Ir and RuCl-cymene. *Dalton Trans.* **2003**, 4132–4138. (b) Li, L.; Brennessel, W. W.; Jones, W. D. C-H Activation of Phenyl Imines and 2-Phenylpyridines with [Cp*MCl₂]₂ (M = Ir, Rh): Regioselectivity, Kinetics, and Mechanism. *Organometallics* **2009**, *28*, 3492–3500.
- (10) For recent reviews on metal-catalyzed asymmetric transfer hydrogenation, see: (a) Ito, J.; Nishiyama, H. Recent Topics of Transfer Hydrogenation. *Tetrahedron Lett.* **2014**, *55*, 3133–3146. (b) Foubelo, F.; Nájera, C.; Yus, M. Catalytic Asymmetric Transfer Hydrogenation of Ketones: Recent Advances. *Tetrahedron: Asymmetry* **2015**, *26*, 769–790. (c) Wang, D.; Astruc, D. The Golden Age of Transfer Hydrogenation. *Chem. Rev.* **2015**, *115*, 6621–6686. (d) Foubelo, F.; Yus, M. Catalytic Asymmetric Transfer Hydrogenation of Imines: Recent Advances. *Chem. Rev.* **2015**, *15*, 907–924. (e) Werkmeister, S.; Neumann, J.; Junge, K.; Beller, M. Pincer-Type Complexes for Catalytic (De)Hydrogenation and Transfer (De)Hydrogenation Reactions: Recent Progress. *Chem. - Eur. J.* **2015**, *21*, 12226–12250. (f) Li, Y. Y.; Yu, S. L.; Shen, W. Y.; Gao, J. X. Iron-, Cobalt-, and Nickel-Catalyzed Asymmetric Transfer Hydrogenation and Asymmetric Hydrogenation of Ketones. *Acc. Chem. Res.* **2015**, *48*, 2587–2598. (g) Chelucci, G.; Baldino, S.; Baratta, W. Ruthenium and Osmium Complexes Containing 2-(aminomethyl)pyridine (Ampy)-based Ligands in Catalysis. *Coord. Chem. Rev.* **2015**, *300*, 29–85. (h) Nedden, H. G.; Zanolli-Gerosa, A.; Wills, M. *Chem. Rev.* **2016**, *16*, 467

- 468 2623–2643. (i) Ayad, T.; Phansavath, P.; Ratovelomanana-Vidal, V.
469 Transition - Metal - Catalyzed Asymmetric Hydrogenation and
470 Transfer Hydrogenation: Sustainable Chemistry to Access Bioactive
471 Molecules. *Chem. Rec.* **2016**, *16*, 2754–2771. (j) Wu, X.; Wang, C.;
472 Xiao, J. Transfer Hydrogenation in Water. *Chem. Rec.* **2016**, *16*,
473 2772–2786. (k) Echeverria, P. G.; Ayad, T.; Phansavath, P.;
474 Ratovelomanana-Vidal, V. Recent Developments in Asymmetric
475 Hydrogenation and Transfer Hydrogenation of Ketones and Imines
476 through Dynamic Kinetic Resolution. *Synthesis* **2016**, *48*, 2523–2539.
477 (l) Štefane, B.; Požgan, F. Metal-Catalysed Transfer Hydrogenation of
478 Ketones. *Top. Curr. Chem.* **2016**, *374*, 1–67. (m) Wills, M. Imino
479 Transfer Hydrogenation Reductions. *Top. Curr. Chem.* **2016**, *374*,
480 69–104. (n) Milner, L.; Talavera, G.; Nedden, H. G. Transfer
481 Hydrogenation Catalysis of Ketones and Imines. *Chim. Oggi* **2017**, *35*,
482 37–40. (o) Mezzetti, A. Iron Complexes with Chiral N/P
483 Macrocycles as Catalysts for Asymmetric Transfer Hydrogenation.
484 *Isr. J. Chem.* **2017**, *57*, 1090–1105.
- 485 (11) Zhou, X.; Wu, X.-F.; Yang, B.-L.; Xiao, J. Varying the Ratio of
486 Formic Acid to Triethylamine Impacts on Asymmetric Transfer
487 Hydrogenation of Ketones. *J. Mol. Catal. A: Chem.* **2012**, *357*, 133–
488 140.
- 489 (12) (a) Xu, Z.; Zhu, S.; Liu, Y.; He, L.; Geng, Z.; Zhang, Y.
490 Synthesis of Optically Active β -Amino Alcohols by Asymmetric
491 Transfer Hydrogenation of α -Amino Ketones. *Synthesis* **2010**, *2010*,
492 811–817. (b) Kumaraswamy, G.; Ramakrishna, G.; Raju, R.;
493 Padmaja, M. An Expedient Synthesis of Enantioenriched Substituted
494 (2-benzofuryl)arylcarbinols via Tandem Rap–Stoermer and Asym-
495 metric Transfer Hydrogenation Reactions. *Tetrahedron* **2010**, *66*,
496 9814–9818. (c) Soni, R.; Jolley, K. E.; Clarkson, G. J.; Wills, M.
497 Direct Formation of Tethered Ru(II) Catalysts Using Arene
498 Exchange. *Org. Lett.* **2013**, *15*, 5110–5113. (d) Soni, R.; Hall, T.
499 H.; Mitchell, B. P.; Owen, M. R.; Wills, M. Asymmetric Reduction of
500 Electron-Rich Ketones with Tethered Ru(II)/TsDPEN Catalysts
501 Using Formic Acid/Triethylamine or Aqueous Sodium Formate. *J.*
502 *Org. Chem.* **2015**, *80*, 6784–6793. (e) Kišic, A.; Stephan, M.; Mohar,
503 B. ansa-Ruthenium(II) Complexes of R₂NSO₂DPEN-(CH₂)_n(η^6 -
504 Aryl) Conjugate Ligands for Asymmetric Transfer Hydrogenation of
505 Aryl Ketones. *Adv. Synth. Catal.* **2015**, *357*, 2540–2546. (f) Touge,
506 T.; Nara, H.; Fujiwhara, M.; Kayaki, Y.; Ikariya, T. Efficient Access to
507 Chiral Benzhydrols via Asymmetric Transfer Hydrogenation of
508 Unsymmetrical Benzophenones with Bifunctional Oxo-Tethered
509 Ruthenium Catalysts. *J. Am. Chem. Soc.* **2016**, *138*, 10084–10087.
510 (g) Cotman, A. E.; Cahard, D.; Mohar, B. Stereoarrayed CF₃-
511 Substituted 1,3-Diols by Dynamic Kinetic Resolution: Ruthenium(II)
512 - Catalyzed Asymmetric Transfer Hydrogenation. *Angew. Chem., Int.*
513 *Ed.* **2016**, *55*, 5294–5298.
- 514 (13) Chen, H.-Y. T.; Wang, C.; Wu, X.; Jiang, X.; Catlow, C. R. A.;
515 Xiao, J. Iridicyclic - Catalysed Imine Reduction: An Experimental and
516 Computational Study of the Mechanism. *Chem.—Eur. J.* **2015**, *21*,
517 16564–16577.
- 518 (14) Jeffrey, G. A. *An Introduction to Hydrogen Bonding*; Oxford
519 University Press: New York, 1997.
- 520 (15) (a) Noyori, R.; Hashiguchi, S. Asymmetric Transfer Hydro-
521 genation Catalyzed by Chiral Ruthenium Complexes. *Acc. Chem. Res.*
522 **1997**, *30*, 97–102. (b) Ikariya, T.; Murata, K.; Noyori, R. Bifunctional
523 Transition Metal-based Molecular Catalysts for Asymmetric Syn-
524 theses. *Org. Biomol. Chem.* **2006**, *4*, 393–406. (c) Ikariya, T.; Blacker,
525 A. J. Asymmetric Transfer Hydrogenation of Ketones with Bifunc-
526 tional Transition Metal - Based Molecular Catalysts. *Acc. Chem. Res.*
527 **2007**, *40*, 1300–1308.
- 528 (16) Yu, J. F.; Long, J.; Yang, Y. H.; Wu, W. L.; Xue, P.; Chung, L.
529 W.; Dong, X. Q.; Zhang, X. M. Iridium-Catalyzed Asymmetric
530 Hydrogenation of Ketones with Accessible and Modular Ferrocene-
531 Based Amino-phosphine Acid (f-Ampha) Ligands. *Org. Lett.* **2017**, *19*,
532 690–693.